Georgia Journal of Science

Volume 71 No. 2 Scholarly Contributions from the Membership and Others

Article 6

2013

Leslie Matrices and Women Population in the United States of America

Brittney Nelson

Denise T. Reid

Antonija Tangar

Jose A. Velez-Marulanda Valdosta State University, javelezmarulanda@valdosta.edu

Follow this and additional works at: https://digitalcommons.gaacademy.org/gjs



Part of the Mathematics Commons

Recommended Citation

Nelson, Brittney; Reid, Denise T.; Tangar, Antonija; and Velez-Marulanda, Jose A. (2013) "Leslie Matrices and Women Population in the United States of America," Georgia Journal of Science, Vol. 71, No. 2, Article 6. Available at: https://digitalcommons.gaacademy.org/gjs/vol71/iss2/6

This Research Articles is brought to you for free and open access by Digital Commons @ the Georgia Academy of Science. It has been accepted for inclusion in Georgia Journal of Science by an authorized editor of Digital Commons @ the Georgia Academy of Science.

LESLIE MATRICES AND WOMEN POPULATION IN THE UNITED STATES OF AMERICA

Brittney Nelson, Denise T. Reid, Antonija Tangar and José A. Vélez-Marulanda* Valdosta State University, Valdosta, GA 31698 *Corresponding author javelezmarulanda@valdosta.edu Fax: (229) 219-1257

ABSTRACT

This research tests the accuracy of the Leslie matrix, which is a discrete age-structured method that uses fertility and survival rates, as a tool for predicting women population. Based on available data for the year 2000, we have constructed a Leslie matrix that predicts female population in the United States for every five years from the years 2000 to 2020. To test the accuracy of this method, we compare the aforementioned obtained projected data for the year 2010 with the actual data for women population in the United States obtained by the 2010 U.S. Census.

Key words: Leslie matrices, population.

INTRODUCTION

In "Essay on the Principle of Population," Thomas Malthus was the first one to offer a scientific explanation and variables influencing long-term population projections (1). Such pioneering work prompted passing of Census Act 1800, the first national census in Britain that has been collected every decade since. In any projection, dimensions of population are the first to be considered. The first complex dimension accounts for age and sex of the species observed. In demography, a key dimension of population dynamics is the distinction between men and women. To clarify, the population growth almost completely correlates to the fertility rates of females. Thus, just by observing female population and the growth within this subgroup, it is possible to predict population growth.

METHODS AND DATA

In 1945, P. H. Leslie constructed a matrix as a means to project population of different species. The Leslie matrix requires three different categories of data divided into equal age groups: survival rates for each age group of females s_i , fertility rates for each age group of female population F_i , and the initial female population for each age group p_i . The matrix is of dimension $n \times n$, where n is number of age groups. Data is distributed and calculated as follows: fertility data occupies the first row of the matrix while survival rates occupy the first sub-diagonal. To project population data, the t-th power of the Leslie matrix is multiplied by the initial population vector x_0 , where t is the number of years between the initial and final year, i.e.,

1

$$x_t = L^t x_0$$

where L is the matrix

$$L = \begin{pmatrix} F_0 & F_1 & F_2 & \dots & F_{n-2} & F_{n-1} \\ s_0 & 0 & 0 & \dots & 0 & 0 \\ 0 & s_1 & 0 & \dots & 0 & 0 \\ 0 & 0 & s_2 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & s_{n-2} & 0 \end{pmatrix}$$

For all $1 \le i \le n$, let $l_i = s_0 s_1 \dots s_{i,1}$ and $m_i = F_{i,1}/s_{i,1}$, i.e., l_i is the fraction of females surviving from birth to age i and m_i is the number of females born, on average, to a female of age i. If λ is a complex eigenvalue of the matrix L, then λ satisfies the discrete Euler-Lotka equation

$$1 = \sum_{i=1}^{n} \lambda^{-i} l_i m_i.$$

For further information, see (2), (3) and (4).

We used Census-based data for the year 2000 (see (5)) and applied the Leslie matrix model to project population for the next two consecutive 5-year intervals by using fertility and survival rates from (6). To compare the effectiveness of the Leslie matrix, we also used simulated data provided in (7), which uses previous projections and modified data to predict population. Moreover, we also analyze the spectrum (i.e., the set of eigenvalues) of each of the Leslie matrices obtained by using both sets of data, with the goal to compare our results with the conclusions of the Perron-Frobenius Theorem (see e.g. (3)), which claims that any matrix with nonnegative real entries has a unique largest real eigenvalue. The Leslie matrices, their eigenvalues and the predictions for each set of data were obtained by using the software MATHEMATICA 8.

RESULTS

Projections for the year 2010 obtained by using the Leslie matrix constructed from fertility and survival rates in Table I versus the data provided for female population in (5) is shown in Figure 1. To better compare actual versus projected values, Figure 2 shows how unreliable the data is by presenting the percent error per age group. In comparison, simulated data published in (7) gives better results, as shown in Table II. This data has been modified to fit the anticipated projections and thus result in better accuracy. In Figures 3 and 4, the complex eigenvalues of the Leslie Matrices corresponding to each set of data are shown. We can see that the Leslie matrices corresponding to the data in Table I and Table II, respectively, have a real eigenvalue that is larger than the others, which satisfies the conclusion of the Perron-Frobenius Theorem. However, the magnitudes of the eigenvalues corresponding to

the Leslie matrix obtained from Table II are very similar as shown in Figure 4, whereas the eigenvalues of the Leslie matrix obtained from Table I are significatively different as shown in Figure 3.

DISCUSSION

Based on our results, we believe Leslie matrix projections are not reliable for the age groups 0-10 and age groups over 85 years old, but are reliable for the rest of the age groups. This claim is supported by our percent error calculations, which show maximum error in projection to be 8%, whereas at extremes it exceeds well above 50%. Furthermore, by the behavior of eigenvalues, we can see that available data is not enough to accurately predict the population. In these age categories, we believe this is caused by inaccurate data in fertility rates in the 0-10 and over 60 age groups. While the rates may be very small, they are far from insignificant and result in such extreme deviations of projections from accurate data. Simulated data used in (7), give more accurate projections, which leads to conclusion that not all data is true and/or available for age groups under 10 years old. When data for these age groups is manipulated, the results are closer to the true value, which is confirmed the data in Table II. Moreover, we have concluded that in order to give better predictions using manipulated data, the available information has to be manipulated in such a way that the corresponding Leslie matrix has complex eigenvalues whose magnitudes are very close to the magnitude of the unique largest real eigenvalue.




Figure 1. Comparison between actual vs. projected values.




Figure 2. Projection error between actual vs. projected values.

Table I. Results using data from (6).

			Actual Population		Estimated	Estimated Population		Actual Population
Age Group	Fertility rates F_i	Survival rates s _i	2002	2005	2010	2015	2020	2010
5	0	0.993164	9.36507	4.39275	4.19878	4.11921	4.13708	9.88194
10	0.000001	0.998622	10.0262	9.30105	4.36273	4.17008	4.09105	9.95902
15	90000.0	23666.0	10.0079	10.0124	9.28823	4.35671	4.16433	10.0973
20	6000'0	608866.0	9.82889	10.0006	10.0051	9.28145	4.35353	10.7367
25	0.0485	0.996493	9.27619	9.81221	9:8836	9.98812	9.2657	10.5718
30	0.1123	0.997464	9.58258	9.24366	6.77779	9.94859	9.9531	10.4663
35	0.1214	918966.0	10.1886	9.55827	9.22021	9.753	9.92336	9:3656
40	0.0941	0.995024	11.388	10.1517	9.52363	9.18679	9.71764	10.1376
45	0.0404	0.993177	11.3128	11.3313	10.1012	9.47624	9.14108	10.497
50	0.0079	0.990092	10.2029	11.2356	11.254	10.0323	9.41158	11.4995
22	9000.0	0.984612	8.97782	10.1018	11.1243	11.1425	9.93285	11.3649
09	0.0005	0.975703	6.96051	8.83967	9.94636	10.9531	10.971	10.1412
99	0.0003	0.960272	5.66882	6.79139	8.6249	9.70469	10.6869	8.74042
70	0.0002	0.936197	5.13318	5.44361	6.52158	8.28225	9.31915	6.58272

Table I. (Continued)

5.03419	4.13541	3.44895	2.34659	1.02398	0.288981	0.044202
7.75381	5.52847	3.96956	2.95407	2.0762	1.08771	0.040397 0.105737 0.206406 0.324282 0.416046 0.044202
6.10548	4.61466	3.74319	3.04558	2.02301	0.899747	0.324282
5.09629	4.3515	3.85914	2.96755	1.67342	0.446378 0.701297 0.899747	0.206406
4.80567	4.48629	3.76027	2.45474	1.30432	0.446378	0.105737
4.95453	4.37136	3.11047	1.91332	0.830206	0.228669	0.040397
0.905493	0.860207	0.789186	0.681708	0.537671	0.462403	0.236645
0.0001	0.00001	0	0	0	0	0
75	80	85	06	92	100	105

Table II. Results using data from (7).

			Actual Population		Estimated	Estimated Population		Actual Population
Age F Group	Fertility rates $F_{_i}$	Survival rates s _i	2002	2002	2010	2015	2020	2010
5	0	0.993164	9.36507	9.15632	9.59477	9.61736	9.02188	9.88194
10	0	0.998622	10.0262	9.30105	9.09373	9.52918	9.55161	9.95902
15	0.0009	0.99927	10.0079	10.0124	9.28823	9.0812	9.51605	10.0973
20	0.0477	0.998303	9.82889	10.0006	10.0006 10.0051	9.28145	9.07457	10.7367

Table II. (Continued)

998	9.58258 9.24366
327	10.1886 9.55827
117	11.388 10.1517
13	11.3128 11.3313
56	10.2029 11.2356
18	8.97782 10.1018
67	6.96051 8.83967
39	5.66882 6.79139
51	5.13318 5.44361
27	4.95453 4.80567
29	4.37136 4.48629
27	3.11047 3.76027
74	1.91332 2.45474
32	0.830206 1.30432
78	0.228669 0.446378
37	0.040397 0.105737




Figure 3. Eigenvalues of the Leslie matrix corresponding to the fertility and survival rates from Table I.




Figure 4. Eigenvalues of the Leslie matrix corresponding to the fertility and survival rates from Table II.

REFERENCES

- 1. Lutz W and Samir K: Dimensions of global population projections: What do we know about future population trends and structures? Philos. Trans. R. Soc. Lond. B Biol. Sci., 365(1554): 2779–2791, 2010. URL: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_14.pdf [cited August 21, 2012].
- 2. Caswell H: "Matrix Population Models: Construction, Analysis, and Interpretation." Sinauer Associates, 2nd edition, 2001.
- 3. Kot M: "Elements of Mathematical Ecology." Cambridge University Press, p377, p386, 2001.
- 4. Leon S J: "Linear Algebra with Applications." Prentice Hall, 8th edition, 2010.
- 5. Interim State Population Projections, 2005. Technical report, U.S. Census Bureau, Population Division, January 2012. URL: http://www.census.gov/population/www/projections/files/Summary-TabC3.pdf [cited August 21, 2012].
- Sandy S F and Bindra K: Vital statistics of California, Table 1-5 2003. URL: http://s3.amazonaws.com/zanran_storage/www.cdph. ca.gov/ ContentPages/34385493.pdf#page=20 [cited August 21, 2012].
- Child Trends Data Bank: Birth and fertility birth and fertility rates. Technical report, March 2012. URL: http://www.childtrendsdata-bank.org/sites/default/files/79_Birth_Rate.pdf [cited August 21, 2012].