

2019


A Tentative List of the Land Snails of Georgia, U.S.A.

Zachary I. Felix
Reinhardt University, zif@reinhardt.edu

Michael A. Dubuc
Reinhardt University

Hassan A. Rana
Reinhardt University

Follow this and additional works at: <https://digitalcommons.gaacademy.org/gjs>

 Part of the [Biodiversity Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Recommended Citation

Felix, Zachary I.; Dubuc, Michael A.; and Rana, Hassan A. (2019) "A Tentative List of the Land Snails of Georgia, U.S.A.," *Georgia Journal of Science*, Vol. 77, No. 2, Article 8.

Available at: <https://digitalcommons.gaacademy.org/gjs/vol77/iss2/8>

This Research Articles is brought to you for free and open access by Digital Commons @ the Georgia Academy of Science. It has been accepted for inclusion in Georgia Journal of Science by an authorized editor of Digital Commons @ the Georgia Academy of Science.

A Tentative List of the Land Snails of Georgia, U.S.A.

Acknowledgements

We thank Shayla Scott for help with building our database. Thanks to the following individuals for sharing museum data: Adam Baldinger, Clarissa Bey, Rudiger Bieler, Cheryl Bright, Brian Helms, Christine Johnson, Timothy Pearce, Gary Rosenburg, Leslie Skibinski, John Slapcinsky, Jamie Smith, and Lee Taehwan. Timothy Pearce, Kathryn Perez, Amy VanDevender, Wayne VanDevender and John Slapcinsky helped tremendously with sorting out taxonomic issues. Helpful reviews were provided by the VanDevenders as well as John Slapcinsky.

A TENTATIVE LIST OF THE LAND SNAILS OF GEORGIA, U.S.A.

Zach I. Felix¹, Michael Dubuc, and Hassan Rana
Biology Program at Reinhardt University
¹corresponding author: zif@reinhardt.edu

ABSTRACT

Because of their high ecological and conservation value, and because we know so little about the group, we compiled a systematic if tentative list of land snails from the state of Georgia. After gleaning a list of species from a monograph on the land snails of eastern United States, written by Leslie Hubricht in 1985, we realized that many species whose ecological requirements are found in Georgia had not been documented there. Therefore, we developed a qualitative model to predict the likelihood that these candidate species occur in Georgia and would eventually be documented. We tested the model with collections data from nine natural history museums and found that the model nonrandomly predicted the species that were collected after the publication of Hubricht's work. Our searches revealed 214 species of land snails collected in Georgia that exist in museums. Our model predicted that another 68 species are likely occur in the state and await documentation. There are at least 10 species of exotic snails within Georgia's borders, some of them invasive. We consider our list of land snails in Georgia tentative but useful because of our systematic approach. It is our hope that more researchers will consider Georgia land snails as a model for studying systematics, evolution, ecology, and conservation.

INTRODUCTION

Land snails, because of their high density and diversity, exhibit high ecological and conservation value. Land snails are a major prey item for numerous vertebrate species (Beissinger 1983; Gunzberger 1999), and can also serve as a major consumer of forest leaf litter (Mason 1974). Snails, because of their calcium-rich shells, are an important food source for passerine birds and, when snails decline, these birds can experience egg defects such as thin or porous shells (Graveland 1996; Graveland et al. 1994). Terrestrial mollusks also serve as food for organisms as diverse as platyhelminths, firefly larvae, and carabid beetles (Barker 2004). By transforming consumed lichens as well as limestone bedrock into feces, snails contribute significantly (>10% of total soil nitrogen) to nutrient cycles of some ecosystems (Jones and Shachack 1990). The diversity and abundance of land snails are positively related to levels of soil calcium in central Appalachians forests (Hotopp 2002). Snails are important dispersal agents for the seeds of certain plants (Turke et al. 2010) and can improve germination of ingested seeds (Calvino-Cancela and Rubido-Bara 2012). When introduced outside of their native range, land snails can cause extensive damage to plants, including rare species (Joe and Daehler 2008), to native animals, especially native land snails (Civeyrel and Simberloff 1996; Meyer and Cowie 2010), as well as economic damage to agricultural systems (Civeyrel and Simberloff 1996).

Globally, land snails, including slugs, reach high levels of diversity within a given ecosystem (e.g., De Winter and Gittenberger 1998; Douglas et al. 2014; Schilthuizen and Rutjes 2001), and the southern Appalachian Mountains are considered a diversity hotspot

for land snails based on overall species richness and diversity as well as on the large numbers of endemic species (Solem 1984). For example, we know a great deal about the biodiversity of the Great Smoky Mountains National Park thanks to the All Taxa Biodiversity Inventory (ATBI), which took place there (Nichols and Langdon 2007). The results of the ATBI (Dourson et al. 2013) revealed that the park contains more species (146) of snails and slugs than species of amphibians, reptiles, mammals, crayfishes, odonates, tardigrades, or orthopterans (Discover Life in America 2016). Of the 18 species of land snails listed as either *threatened* or *endangered* by the United States Fish and Wildlife Service, 10 species are found in the lower 48 states, and only five are found in the southeastern United States (USFWS 2015).

Considering these values, it is surprising how little is known about the land snail fauna of the Southeast. Georgia remains terra incognita in terms of its land snail inhabitants. The first published list of land snail species in the state of Georgia was by Hubricht (1964). Even at that time, the author remarked that less was known about the land snails in Georgia than almost any other place in the eastern United States. The list included a total of 123 species personally collected by the author. In his monograph on the distribution of eastern land snails, Hubricht (1985) listed a total of 150 species from within Georgia's boundaries. Since this time, research on Georgia land snails has focused almost exclusively on cave ecosystems. One study reported three land snail families containing nine species from caves in Georgia (Holsinger and Peck 1971), while another reported at least nine species from two families (Reeves et al. 2000). A review of the obligate cave fauna of Georgia listed two species of land snails—*Glyphyalinia specus* and *Helicodiscus barri* (Niemiller et al. 2012). Because of this knowledge gap, we summarized data on land snails collected in Georgia in order to compile a list of recorded and likely species for the state.

For this paper, we used an approach that combined the use of literature with a search of museum records to compile a list of terrestrial snail species, including native and exotic species, for which a vouchered specimen collected from the state of Georgia exists. We also developed and tested a qualitative model to predict species that have not been found in Georgia yet, but are likely occur here based on the proximity of known populations and ecoregional affinities. The same model was used to identify species that have vouchered Georgia specimens, but that we are doubtful occur in the state. This effort is the type of basic research needed for us to properly understand regional patterns in land snail diversity, and for managers to pinpoint locations for conservation efforts (Lydeard et al. 2004).

MATERIALS & METHODS

Our first step was to compile a list of the species shown in Hubricht (1985) as occurring in at least one county in the state of Georgia. This monograph remains the most extensive work on land snail distributions in the eastern United States to date. One impression that we gained during this process was that the state of Georgia was not well-represented in the sampling, and that many species whose ecological requirements should be met within Georgia ecosystems were not recorded there as of the time of Hubricht's 1985 publication. Therefore, we constructed a qualitative model to predict which species might eventually be recorded in Georgia based on their geographic proximity and level III ecoregional associations (US EPA 2006). Using this model we classified every species not recorded by Hubricht 1985 in any Georgia counties as likely, possible, or unlikely to actually occur in

Georgia. The following are the rules of the model: If a species is not found in any adjacent states, and Georgia is not included in gaps between known sites, then the species was classified as *unlikely*. If the species is found in one adjacent state, but not within three counties of Georgia (excluding species restricted to Florida Keys); or is found in two or more surrounding states (one or more being nonadjacent), and Georgia is in the gap between known sites; or is found in two or more adjacent states in ecoregions not found in Georgia, and Georgia is in gap between these states; then the species is classified as *possible*. If, on the other hand, the species is found in one or more adjacent states within three counties of Georgia; or is found in two or more adjacent states in ecoregions found in Georgia, and Georgia is in a gap between these states; then the species was classified as *likely*.

On 01/23/12, a request was sent to the following museums for all digital records from the state of Georgia for all families of land snails included in Hubricht 1985: United States National Museum, Washington (USNM); Academy of Natural Sciences Museum, Philadelphia (ANSP); Museum of Comparative Zoology, Cambridge (MCZ); Field Museum of Natural History, Chicago (FMNH); University of Michigan Museum of Zoology, Ann Arbor (UMMZ); Carnegie Museum of Natural History (CMNH); Auburn University Museum of Natural History, Auburn (AUM); Delaware Museum of Natural History, Wilmington (DMNH); Florida Museum of Natural History, Gainesville (UF); American Museum of Natural History, New York (AMNH); Ohio State University Museum of Biological Diversity (OSU); and North Carolina State Museum of Natural History (NCSM). These collections include all of those listed used by both Hubricht (1985) and Minton and Perez (2010). All records received were compiled into a common spreadsheet and taxonomic names were all updated for consistency. Taxonomy followed Turgeon et al. 1988 at the species level in most cases. Exceptions were accounted for by literature such as Emberton (1988, 1991). Taxonomy followed Bouchet et al. 2017 at the family level with some exceptions; we chose to follow Hausdorf (1998) and Slapcinsky (2018) in assignment of the genera *Glyphyalinia*, *Nesovitrea*, *Paravitrea*, *Pilsbryna*, and *Mesomphix* to Oxychilidae.

We used museum data, particularly those collected after the publication of Hubricht (1985), to test the validity of our predictive model. All species listed in Hubricht (1985), but not reported in Georgia by the same, were considered as candidate species for eventual discovery within the state. We used Chi-square to test the null hypothesis that, for the species documented in Georgia in museum collections but not in Hubricht (1985), the proportion found in the different likelihood categories is random. For random, or expected, values we used the proportions of all 372 candidate species found in the three likelihood categories. If, alternatively, our model was useful, then more candidate species that our model classified as likely to occur in Georgia would be found in museum collections than expected at random, and, on the other hand, fewer candidate species that our model classified as unlikely would be found in museum collections than expected at random.

RESULTS

A total of 5232 records were returned from each of these respective collections: AMNH (314), ANSP (395), AUM (33), CMNH (311), DMNH (157), FMNH (2723), UF (1000), MCZ (282), NCSM (18). From these records, we found a total of 214 unique species that were collected in the state of Georgia from 27 families and 63 genera. There were three

species listed in Hubricht (1985) that were not represented in the museums that we surveyed, thus bringing the total number of land snail species recorded in Georgia to 217 (Table I).

We found that 64 species uncovered in our survey of museum specimens were not listed as occurring in Georgia by Hubricht (1985). Twenty-three of these species were not found in Hubricht at all and represent either exotic species not considered by Hubricht in his work or species that have been described since the publication. Our model nonrandomly predicted the 41 other species that were found in Georgia after the publication Hubricht (1985) from the list of 372 candidate species not recorded in Georgia by Hubricht ($P < 0.001$, $X^2 = 41.95$, $df = 2$). Only 101 of the 372 (27%) of the candidate species were predicted as *likely*, whereas 29 of the 41 species subsequently found in Georgia (71%) were predicted as *likely* by our model. On the other hand, 200 of the 372 candidate species not recorded in Georgia by Hubricht (54%) were predicted as *unlikely* by our model, whereas only five (1.3%) of the species recorded in museums additional to Hubricht were those that were predicted as *unlikely*. For those species predicted as *possible*, there were 71/372 candidates (19%) and 7/41 (17%) subsequently recorded. These results lend some support to our predictions of which species will eventually be found in Georgia. After considering the species found in our museum survey, this leaves 69 species of snails that our model classified as likely to occur in the state but have still never been documented (Table II).

The following 12 species of exotic snails have been documented in the state of Georgia (Table I): *Allopeas clavulinum*, *Allopeas gracile*, *Arion circumscriptus*, *Cochlicopa lubrica*, *Hemicycla plicaria*, *Limax maximus*, *Opeas pumilum*, *Opeas pyrgula*, *Otala lactea*, *Otala punctata*, *Oxychilus draparnaudi*, *Rumina decollata*, *Subulina octona*.

Table I. Species of land snails collected in Georgia and the minimum number of specimens found in various museum collections. Specimen numbers are minimums in some cases because if museum records did not contain data on specimen numbers we recorded “ ≥ 1 ” for this field. AMNH = American Museum of Natural History, ANSP = Academy of Natural Sciences of Philadelphia, AUM = Auburn Museum of Natural History, CMNH = Carnegie Museum of Natural History, DMNH = Delaware Museum of Natural History, FMNH = Field Museum, UF = Florida Museum of Natural History, MCZ = Museum of Comparative Zoology at Harvard, NCSM = North Carolina State Museum.

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Achatinidae									
<i>Allopeas clavulinum</i> ^e						≥ 1			
<i>Allopeas gracile</i> ^e		10				46			
<i>Opeas pumilum</i> ^e							1		
<i>Opeas pyrgula</i> ^e						31			
<i>Rumina decollata</i> ^e		18		57		112		12	
<i>Subulina octona</i> ^e								2	
Agriolimacidae									
<i>Deroceras laeve</i>	7						3		
Arionidae									
<i>Arion circumscriptus</i> ^e						≥ 1			

^e denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Cochlicopidae									
<i>Cochlicopa lubrica</i> ^e			16						
<i>Cochlicopa morseana</i>						9			
Discidae									
<i>Anguispira alternata</i>		94		42	1	5		110	
<i>Anguispira fergusonii</i>						101			
<i>Anguispira mordax</i>				1					
<i>Anguispira strongylodes</i>						265	23		
<i>Discus patulus</i>		88		57	18	854	7	45	
<i>Discus whitneyi</i>				≥1					
Ellobiidae									
<i>Carychium clappi</i>						217			
<i>Carychium exiguum</i>						99			
<i>Carychium exile</i>						481			
<i>Carychium mexicanum</i>	3					1026			
<i>Carychium nannodes</i>						110			
Euconulidae									
<i>Euconulus chersinus</i>	4	23				280	1		
<i>Euconulus dentatus</i>						3			
<i>Euconulus trochulus</i>	5					33			
<i>Guppya sterkii</i>						4			
Gastrocoptidae									
<i>Gastrocopta armifera</i>				5		111			
<i>Gastrocopta clappi</i>						115			
<i>Gastrocopta contracta</i>	1	1				168			
<i>Gastrocopta corticaria</i>		7				1			
<i>Gastrocopta pellucida</i>	42	4		12		57			
<i>Gastrocopta pentodon</i>	10	16		11		57			
<i>Gastrocopta procera</i>				2		16			
<i>Gastrocopta rupicola</i>	18	30		9		159		3	
<i>Gastrocopta tappaniana</i>	7			1		138			
Gastrodontiidae									
<i>Gastrodonta interna</i>		115	30	92		544	47	123	4
<i>Striatura meridionalis</i>	13		2	1		183			
<i>Ventridens acerra</i>		11		53		94	15	17	
<i>Ventridens arcellus</i>						4			
<i>Ventridens cerinoideus</i>	136	77				1006	6	21	
<i>Ventridens collisella</i>						3			
<i>Ventridens decussatus</i>						17			
<i>Ventridens demissus</i>		13	2			333			
<i>Ventridens gularis</i>		88	2	117		500		202	
<i>Ventridens intertextus</i>		14	1	13		301	6	17	
Gastrodontiidae									
<i>Ventridens lawae</i>						436			
<i>Ventridens ligera</i>		5		1			7	2	1

^e denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Gastrodontidae, continued									
<i>Ventridens pilsbryi</i>						1275	32		
<i>Ventridens suppressus</i>		5				7			
<i>Ventridens theloides</i>						936			
<i>Vitrinizonites latissimus</i>						8			
<i>Zonitoides arboreus</i>	137	138		86		615		12	2
<i>Zonitoides elliotti</i>		25	1	98		98	17	107	
<i>Zonitoides nitidus</i>				12					
<i>Zonitoides patuloides</i>						1			
Haplotrematidae									
<i>Haplotrema concavum</i>		8		10	4	96	3	7	10
Helicidae									
<i>Hemicycla plicaria</i> ^e					8				
<i>Otala lactea</i> ^e					2				
<i>Otala punctata</i> ^e		16				345			
Helicinidae									
<i>Helicina orbiculata</i>	23	16		18		112			
Helicodiscidae									
<i>Helicodiscus barri</i>						2			
<i>Helicodiscus fimbriatus</i>		10		3		19		12	
<i>Helicodiscus inermis</i>	1					36			
<i>Helicodiscus notius</i>						256			
<i>Helicodiscus parallelus</i>	41	25	1	3		643		4	
Limacidae									
<i>Limax maximus</i> ^e						1			
Oxychilidae									
<i>Glyphyalinia carolinensis</i>		9		44	4				
<i>Glyphyalinia cryptomphala</i>		4		3		9			
<i>Glyphyalinia cumberlandiana</i>		2				21	2		
<i>Glyphyalinia indentata</i>		51		19	9	297	4	3	
<i>Glyphyalinia junaluskana</i>						1			
<i>Glyphyalinia luticola</i>						17			
<i>Glyphyalinia pentadelphia</i>		1		2		1			
<i>Glyphyalinia praecox</i>		2				4	1		
<i>Glyphyalinia rhoadsi</i>						13			
<i>Glyphyalinia sculptilis</i>		24		32		41	4		
<i>Glyphyalinia solida</i>	3					105			
<i>Glyphyalinia specus</i>						7			
<i>Glyphyalinia umbilicata</i>	17					154			
<i>Glyphyalinia wheatleyi</i>				1		206			
<i>Mesomphix andrewsae</i>		7			3		2		

^e denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Oxychilidae, continued									
<i>Mesomphix anurus</i>						5			6
<i>Mesomphix capnodes</i>		8		6	6	54	7	4	2
<i>Mesomphix cupreus</i>		2		7				1	
<i>Mesomphix globosus</i>						488	4		
<i>Mesomphix latior</i>		4			2	32	1		
<i>Mesomphix perlaevis</i>			2	33	1	112	8	23	
<i>Mesomphix pilsbryi</i>		8	1	5	3	59			
<i>Mesomphix rugeli</i>						1			
<i>Mesomphix subplanus</i>						1		4	
<i>Mesomphix vulgatus</i>		51		25	12	50		28	
<i>Nesovitrea dalliana</i>	4					245	4		
<i>Nesovitrea electrina</i>				24					
<i>Oxychilus draparnaudi</i> ^e						2			
<i>Paravitrea amicalola</i>						44			
<i>Paravitrea capsella</i>						99		8	
<i>Paravitrea diana</i>						19			
<i>Paravitrea lamellidens</i>									
<i>Paravitrea multidentata</i>						≥1			
<i>Paravitrea petrophila</i>								50	
<i>Paravitrea placentula</i>				5					
<i>Paravitrea umbilicaris</i>						2			
Philomycidae									
<i>Megapallifera mutabilis</i>							1		
<i>Pallifera fosteri</i>	7								
<i>Pallifera unicolor</i>	1								
<i>Philomycus carolinianus</i>	2							2	
<i>Philomycus togatus</i>							2		
Polygyridae									
<i>Allogona profunda</i>								1	
<i>Appalachina sayana</i>					2				
<i>Daedalochila auriculata</i>							≥1		
<i>Daedalochila auriformis</i>		23		17		12	4	38	
<i>Daedalochila avara</i>				2				2	
<i>Daedalochila delecta</i>							6		
<i>Daedalochila leporina</i>		9				4			
<i>Daedalochila postelliana</i>	2	5		9	6	11		29	
<i>Daedalochila subclausa</i>						3			
<i>Daedalochila uvulifera</i>		1							
<i>Euchemotrema fraternum</i>		2		2	6	6	3		
<i>Euchemotrema leai</i>		2		1	4	7			
<i>Fumonelix archeri</i>						3			
<i>Fumonelix christyi</i>						100			

^e denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Polygyridae, continued									
<i>Fumonelix</i>									
<i>clingmanicus</i>						1			
<i>Fumonelix wheatleyi</i>		1				9		2	
<i>Inflectarius</i>									
<i>approximans</i>		3							
<i>Inflectarius downieanus</i>		1							
<i>Inflectarius edentatus</i>						2			
<i>Inflectarius inflectus</i>		45		56	51	753	30	90	
<i>Inflectarius rugeli</i>		16	1	14	6	198	26	11	
<i>Inflectarius smithi</i>					4				
<i>Lobosculum pustula</i>	64	28		4		233	1	12	
<i>Lobosculum pustuloides</i>		9		8	6	382		33	
<i>Mesodon andrewsae</i>				1		1		4	
<i>Mesodon clausus</i>		1	1			3			
<i>Mesodon normalis</i>		4	1	31	27	196	4	3	
<i>Mesodon thyroidus</i>		39		37	7	260	16	45	
<i>Mesodon zaletus</i>		1		1		7	4		
<i>Millerelix dorfeuilliana</i>				3				1	
<i>Millerelix plicata</i>		27		17	73	140	1	18	
<i>Millerelix troostiana</i>					5				
<i>Neohelix albolabris</i>		70		17	25	11		55	
<i>Neohelix dentifera</i>							≥1		
<i>Neohelix divesta</i>					2				
<i>Neohelix major</i>			2	32		199	2	8	
<i>Neohelix solemi</i>						≥1			
<i>Patera appressa</i>			2	9	10	99		34	
<i>Patera clarki</i>		6		9	5	117	9	3	
<i>Patera laevior</i>						17			
<i>Patera perigrapta</i>		20	8	23	16	213	2	16	14
<i>Polygyra cereolus</i>	15	80		21	2	553	20	85	
<i>Polygyra septemvolva</i>	136					138		6	
<i>Praticolella jejuna</i>	30	7		22	2	7	21	4	
<i>Praticolella lawae</i>		4		5	20	29			
<i>Praticolella mobilians</i>		4		1					
<i>Stenotrema altispira</i>					1				
<i>Stenotrema barbatum</i>						1			
<i>Stenotrema barbigerum</i>		20		28	4	98	10	61	
<i>Stenotrema cohuttense</i>		4	56	39	65	127	3	37	
<i>Stenotrema edwardsi</i>				35		4		15	
<i>Stenotrema exodon</i>		6							
<i>Stenotrema florida</i>						6			
<i>Stenotrema hirsutum</i>		3						17	
<i>Stenotrema</i>									
<i>magnifumosum</i>		15		1	19	212	7		
<i>Stenotrema maxillatum</i>		23	10	3	4	109		19	
<i>Stenotrema spinosum</i>		9		12		19	20	4	1
<i>Stenotrema turbinella</i>						4			

^e denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Polygyridae, continued									
<i>Triodopsis affinis</i>						659			
<i>Triodopsis alabamensis</i>				10		260			
<i>Triodopsis fallax</i>		39		2	81	44		19	
<i>Triodopsis fraudulentata</i>								1	
<i>Triodopsis hopetonensis</i>	179	155		35		1298	5	59	3
<i>Triodopsis juxtidentis</i>		1				6		2	
<i>Triodopsis messana</i>						198			
<i>Triodopsis palustris</i>						61			
<i>Triodopsis tennesseensis</i>			3			2			
<i>Triodopsis tridentata</i>		9		24	1	226	5	23	
<i>Triodopsis vannostrandii</i>		23	8	39		312	1	50	
<i>Triodopsis vulgata</i>					7	9	1		
<i>Xolotrema caroliniense</i>		17		3	17	25	3	3	
<i>Xolotrema denotatum</i>				1	2				
<i>Xolotrema fosteri</i>					6				
<i>Xolotrema obstrictum</i>				6				3	
Pomatiopsidae									
<i>Pomatiopsis lapidaria</i>						323			
Pristilomatidae									
<i>Hawaiiia alachuana</i>	2								
<i>Hawaiiia minuscula</i>	51	51		7		320	10		
Punctidae									
<i>Punctum blandianum</i>						13			
<i>Punctum minutissimum</i>	3					152			
<i>Punctum vitreum</i>		1							
Pupillidae									
<i>Pupoides albilabris</i>				20		246		67	
<i>Pupoides modicus</i>	65	67		82	3	17	15	3	
Spiraxidae									
<i>Euglandina rosea</i>	40	21	2	14	9	35		76	3
Strobilopsidae									
<i>Strobilops aeneus</i>	84			1	4	306			
<i>Strobilops hubbardi</i>	4					1			
<i>Strobilops labyrinthicus</i>		3			2	60			
<i>Strobilops texasianus</i>	10	1				63			
Succineidae									
<i>Catinella oklahomarum</i>						41			
<i>Catinella pugilator</i>						1			
<i>Catinella vermeta</i>		12				128			
<i>Novisuccinea ovalis</i>					37	14		13	
<i>Succinea campestris</i>	46	60		16		232		48	
<i>Succinea indiana</i>		12			2	127			
<i>Succinea unicolor</i>	39					216			
<i>Succinea wilsonii</i>	3	9			4	113	8		

^a denotes species that are considered exotic relative to the state of Georgia.

Table I (continued)

Family and species	AMNH	ANSP	AUM	CMNH	DMNH	FMNH	UF	MCZ	NCSM
Truncatellinidae									
<i>Columella simplex</i>						100			
Valloniidae									
<i>Pupisoma dioscoricola</i>	11					246			
<i>Pupisoma macneilli</i>	5					21			
<i>Pupisoma minus</i>						≥1			
<i>Vallonia excentrica</i>						8			
<i>Vallonia pulchella</i>								14	
Vertiginidae									
<i>Vertigo milium</i>						457			
<i>Vertigo oralis</i>	6	1				220			
<i>Vertigo oscariana</i>						1			
<i>Vertigo ovata</i>				5		12			
<i>Vertigo rugosula</i>		1							
<i>Vertigo teskeyae</i>						101			

^e denotes species that are considered exotic relative to the state of Georgia.

Table II. Species of land snails not found in any of the museum collections that we searched, but which our qualitative model predicts as likely to actually occur in the state of Georgia, USA

Family and species of land snails		
Bulimulidae	<i>Helicodiscus saludensis</i>	<i>Daedalochila hausmani</i>
<i>Drymaeus dormani</i>	Haplotrematidae	<i>Daedalochila peninsulæ</i>
<i>Rabdotus dealbatus</i>	<i>Haplotrema kendeighi</i>	<i>Fumonelix jonesianus</i>
Discidae	Oxychilidae	<i>Fumonelix orestes</i>
<i>Anguispira alabama</i>	<i>Glyphyalinia latebricola</i>	<i>Fumonelix wheatleyi</i>
<i>Anguispira cumberlandiana</i>	<i>Glyphyalinia lewisiana</i>	<i>Inflectarius smithi</i>
<i>Anguispira jessica</i>	<i>Glyphyalinia ocoae</i>	<i>Mesodon elevatus</i>
<i>Anguispira knoxensis</i>	<i>Paravitrea bidens</i>	<i>Mesodon sanus</i>
<i>Discus nigrimontanus</i>	<i>Paravitrea calcicola</i>	<i>Millerelix fatigiata</i>
Euconulidae	<i>Paravitrea clappi</i>	<i>Neohelix alleni</i>
<i>Dryachloa dauca</i>	<i>Paravitrea conecuhensis</i>	<i>Patera sargentiana</i>
<i>Euconulus fulvus</i>	<i>Paravitrea lacteodens</i>	<i>Stenotrema calvescens</i>
<i>Guppya gundlachi</i>	<i>Paravitrea metallacta</i>	<i>Stenotrema deceptum</i>
Gastrocoptidae	<i>Paravitrea pilsbryana</i>	<i>Stenotrema depilatum</i>
<i>Gastrocopta servilis</i>	<i>Paravitrea tantilla</i>	<i>Stenotrema edgarianum</i>
Gastrodontidae	<i>Paravitrea tiara</i>	<i>Stenotrema pilula</i>
<i>Striatura ferrea</i>	<i>Paravitrea toma</i>	Punctidae
<i>Ventridens lasmodon</i>	<i>Paravitrea variabilis</i>	<i>Punctum smithi</i>
<i>Ventridens percallosus</i>	<i>Pilsbryna castanea</i>	Succineidae
<i>Ventridens volusiae</i>	Philomycidae	<i>Catinella hubrichti</i>
Helicinidae	<i>Megapallifera wetherbyi</i>	<i>Oxyloma effusa</i>
<i>Hendersonia occulta</i>	<i>Pallifera hemphilli</i>	<i>Succinea chittenangoensis</i>
Helicodiscidae	<i>Pallifera secreta</i>	<i>Succinea urbana</i>
<i>Helicodiscus aldrichianus</i>	<i>Philomycus sellatus</i>	Valloniidae
<i>Helicodiscus bonamicus</i>	<i>Philomycus venustus</i>	<i>Vallonia perspectiva</i>
<i>Helicodiscus hadenoecus</i>	Polygyridae	Vertiginidae
<i>Helicodiscus hexodon</i>	<i>Appalachina chilhoweensis</i>	<i>Vertigo gouldi</i>

DISCUSSION

We will say up front that this work, with all of its limitations, is a starting point for a complete and exhaustive list of land snails in the state of Georgia. One major limitation of our data is the general dearth of collecting in the state, and another is that we had to rely only on digitized museum records that were unverified. Some museums that provided data for Hubricht (1985), including Ohio State University, University of Michigan Museum of Zoology, and the United States National Museum, could not provide any digitized records for the present investigation. However, because our species list includes information from Hubricht (1985), a study that used data from these same collections, our species list appears to be representative of what is found in museum collections; it is notable that only three species listed as occurring in Georgia by Hubricht (1985) were not represented in the digitized museum records. Land snails are notoriously difficult to identify, and there have been many taxonomic changes in the past within this group (J. Slapcinsky, personal communication). Some species in our list have since been split into multiple taxa, and others are almost certainly misidentified. In fact, we used our qualitative model results to categorize species for which a Georgia specimen can be found in museum collections, but that which we doubt actually occur here. Accordingly, our list of 217 species includes five doubtful species (*Appalachina sayana*, *Millerelix dorfeuilliana*, *Neohelix divesta*, *Triodopsis fraudulenta*, and *Zonitoides nitidus*), for which Georgia seems to fall outside of their native range. Interestingly, all but one of these species, *Millerelix dorfeuilliana*, are represented in only one collection; this suggests that these could be misidentified or mislabeled. Future research should reexamine the identifications and verify or refute the legitimacy of these specimens' provenance.

The number of land snails documented here as occurring in Georgia, 217, is on par with the number of species in surrounding states. Minton and Perez (2010) documented 226 species of terrestrial mollusks in neighboring Alabama in a similar analysis of museum records. At the time of publication, the list of land snails for North and South Carolina included 262 and 102 species, respectively (A. VanDevender, personal communication). The true number of species of land snails in Georgia is almost certainly higher than 217, and our best estimate considering the 69 species that our model categorized as likely to occur in Georgia would be 270–290 species. A high number of land snail species for Georgia is not surprising considering the high ecological diversity found within Georgia's borders (Edwards et al. 2013). Our model may overpredict the presence of some high-elevation southern Appalachian species with restricted ranges but, then again, we predict that our state harbors similar undescribed species.

Many of the exotic taxa, e.g., *Allopeas*, *Otala*, *Deroceras*, *Opeas*, that we uncovered as living in Georgia are common invasives globally (Brodie and Barker 2011; Cowie 2001; LaPierre et al. 2010). In California grasslands, *Otala lactea*, an established nonnative snail, prefers to feed on *Brassica nigrans*, a nonnative plant that co-occurs with *O. lactea* in its native range, suggesting that this exotic snail may slow the progress of an invasive plants (LaPierre et al. 2010); however, this is certainly an exception, and most invasive land snail species almost certainly have an overall detrimental effect on indigenous ecosystems. The *O. lactea* specimens found in our query of museum records may be misidentified *Otala punctata*, a species known to have invaded coastal islands near Savannah, and it seems unlikely that the narrowly endemic *Hemicycla plicaria* records are accurate (J. Slapcinsky, pers. comm.). Considering the potential ecological and

economic impacts that exotic land snails could have in Georgia, we suggest that the list of exotic land snail species presented here can serve as the nucleus of an organized surveillance program for the study of the spread of these species, and as a baseline for detection of newly introduced species.

One of our major goals in this investigation was to spur interest in research on land snails in the state of Georgia. Accordingly, an important finding was the estimation that there are 68 species of land snail that are likely to occur in Georgia but have not been documented (Table II). If all of these species were eventually found in the state, the species list for Georgia would increase by 32% compared to the list from the current study, suggesting that sampling effort for this important taxon has been grossly inadequate till present. Though it is unlikely that all likely species will be discovered in Georgia, the results of our model validation indicate that this list is useful in guiding targeted surveys for additions to the state species list. The list of species likely to occur in Georgia contains both macro- and microscopic snails from a variety of families, but certain genera, e.g., *Paravitrea*, *Fumonelix*, and *Helicodiscus*, contain several species to be targeted. Surveys that are designed to add new species to the state list should take place on the limestone outcroppings and seepages on the Cumberland Plateau or in the Blue Ridge Mountains in the northern part of the state, as many of the undocumented species have been found just north of Georgia in these ecoregions. Swamps, wet prairies, and hammocks in south Georgia would also be promising places for malacological exploration. Because of the small size of many snail species, surveyors would do well to sift through leaf litter and look for shells of these minute creatures (Dourson et al. 2013).

We encourage all ecologists, biogeographers, and systematists to consider adopting this fascinating group as a model for their research. With 20 or more species occurring in the same forest floor habitats here in Georgia, these communities offer tremendous opportunities to study topics like niche partitioning, competition, nutrient cycling, and trophic relationships. The sensitivity of land snails to environmental change (Baur and Baur 1993; Bezemer and Knight 2001), and their small home ranges (Edworthy et al. 2012), combined with Georgia's unique geographic position at the end of the Appalachian Mountains, all make Georgia land snails a perfect model to study the rate and nature of distributional shifts in response to climate change (Fortunato 2016). Work needs to be done for us to achieve a better understanding of the evolutionary relationships among the many land snail taxa, and especially of the taxonomy of these groups. Recent molecular genetic studies suggest that many nominal taxa are not monophyletic and require a new taxonomic scheme that more accurately reflects the evolution of the groups (Perez et al. 2014). Based on this same research it is likely that numerous cryptic species of land snails await description within our borders. Many states have more systematically assessed the conservation status of their land snails and assigned protective status to certain species. Georgia is far behind in this regard, and it is our hope that our preliminary list here can serve as a seed for a larger effort to assess and protect this important natural resource.

ACKNOWLEDGMENTS

We thank Shayla Scott for help with building our database. Thanks to the following individuals for sharing museum data: Adam Baldinger, Clarissa Bey, Rudiger Bieler, Cheryl Bright, Brian Helms, Christine Johnson, Timothy Pearce, Gary Rosenburg, Leslie Skibinski, John Slapcinsky, Jamie Smith, and Lee Taehwan. Timothy Pearce, Kathryn

Perez, Amy VanDevender, Wayne VanDevender and John Slapcinsky helped tremendously with sorting out taxonomic issues. Helpful reviews were provided by the VanDevenders as well as John Slapcinsky.

LITERATURE CITED

- Barker, G.M., ed. 2004. Natural Enemies of Terrestrial Molluscs. CABI Publishing. doi:[10.1079/9780851993195.0000](https://doi.org/10.1079/9780851993195.0000).
- Baur, B. and A. Baur. 1993. Climatic warming due to thermal radiation from an urban area as possible cause for the local extinction of a land snail. *Journal of Applied Ecology*, 30, 333–340. doi:[10.2307/2404635](https://doi.org/10.2307/2404635).
- Beissinger, S.R. 1983. Hunting behavior, prey selection, and energetics of snail kites in Guyana: choice by a specialist. *The Auk*, 100, 84–92.
- Bezemer, T.M. and K.J. Knight. 2001. Unpredictable responses of garden snail (*Helix aspera*) populations to climate change. *Acta Oecologica*, 22, 201–208. doi:[10.1016/S1146-609X\(01\)01116-X](https://doi.org/10.1016/S1146-609X(01)01116-X).
- Bouchet P., J.P. Rocroi, B. Hausdorf, A. Kaim, Y. Kano, A. Nützel, P. Parkhaev, M. and E.E. Strong. 2017. Revised classification, nomenclator and typification of gastropod and monoplacophoran families. *Malacologia*, 61, 1–526. doi:[10.4002/040.061.0201](https://doi.org/10.4002/040.061.0201).
- Brodie, G. and G.M Barker. 2011. Introduced land snails in the Fiji Islands: are there risks involved? *In*: C.R. Veitch and D.R. Towns, eds. *Island Invasives: Eradication and Management*. International Union for the Conservation of Nature, pp. 32–36
- Calvino-Cancela, M. and M. Rubido-Bara. 2012. Effects of seed passage through slugs on germination. *Plant Ecology*, 213, 663–673. doi:[10.1007/s11258-012-0030-8](https://doi.org/10.1007/s11258-012-0030-8).
- Civeyrel, L. and D. Simberloff. 1996. A tale of two snails: is the cure worse than the disease? *Biodiversity and Conservation*, 5, 1231–1252.
- Cowie, R.W. 2001. Invertebrate invasions on Pacific islands and the replacement of unique native faunas: a synthesis of the land and freshwater snails. Contribution no. 2001-001 of Bishop Museum's Pacific biological survey. *Biological Invasions*, 3(2), 119–136.
- Cox, G.W. 1999. *Alien Species in American and Hawaii*. Island Press.
- De Winter, A.J. and E. Gittenberger. 1998. The land snail fauna of a square kilometer patch of rainforest in southwestern Cameroon, high species richness, low abundance and seasonal fluctuations. *Malacologia*, 40, 231–250.
- Douglas, D.A., D.C. Dourson, and R.S. Caldwell. 2014. The land snails of White Oak Sinks, Great Smoky Mountains National Park, Tennessee. *Southeastern Naturalist*, 13, 166–175. doi:[10.1656/058.013.0116](https://doi.org/10.1656/058.013.0116).
- Dourson, D.C., K. Langdon, and J. Dourson. 2013. *Land Snails of the Great Smoky Mountains National Park and Southern Appalachians, Tennessee and North Carolina*. Goatslug Publications.
- Edwards, L., J. Ambrose, and L. K. Kirkman. 2013. *The Natural Communities of Georgia*. The University of Georgia Press.
- Edworthy, A.B., K.M.M. Steensma, H.M. Zandberg, and P.L. Lilley. 2012. Dispersal, home-range size, and habitat use of an endangered land snail, the Oregon forestsnail (*Allogona townsendiana*). *Canadian Journal of Zoology*, 90, 875–884. doi:[10.1139/z2012-056](https://doi.org/10.1139/z2012-056).

- Emberton, K.C. 1988. The genitalic, allozymic, and conchological evolution of the eastern North American Triodiopsinae (Gastropoda: Pulmonata: Polygyridae). *Malacologia*, 28, 159–273.
- Emberton, K.C. 1991. The genitalic, allozymic, and conchological evolution of the Tribe Mesodontini (Pulmonata: Stylommatophora: Polygyridae). *Malacologia*, 33, 71–178.
- Endangered species: U.S. Species. 2015. United States Fish and Wildlife Service. <https://www.fws.gov/endangered/species/us-species.html>. Accessed 19 June 2016.
- Fortunato, H. 2016. Mollusks: tools in environmental and climate research. *American Malacological Bulletin*, 33, 310–324. doi:[10.4003/006.033.0208](https://doi.org/10.4003/006.033.0208).
- Graveland, J., R. van der Wal, J.H. van Balen, and A.J. van Noordwijk. 1994. Poor reproduction in forest passerines from decline of snail abundance on acidified soils. *Nature*, 368, 446–448. doi:[10.1038/368446a0](https://doi.org/10.1038/368446a0).
- Graveland, J.R. 1996. Avian eggshell formation in calcium-rich and calcium-poor habitats: importance of snail shells and anthropogenic calcium sources. *Canadian Journal of Zoology*, 74, 1035–1044. doi:[10.1139/z96-115](https://doi.org/10.1139/z96-115).
- Gunzberger, M.S. 1999. Diet of the Red Hills salamander *Phaeognathus hubrichti*. *Copeia*, 1992, 523–525. doi:[10.2307/1447504](https://doi.org/10.2307/1447504).
- Hausdorf, B. 1998. Phylogeny of the Limacoidea *sensu lato* (Gastropoda: Stylommatophora). *Journal of Molluscan Studies*, 64, 35–66. doi:[10.1093/mollus/64.1.35](https://doi.org/10.1093/mollus/64.1.35).
- Holsinger J.R. and S.B. Peck. 1971. The invertebrate cave fauna of Georgia. *Bulletin of the National Speleological Society*, 33, 23–44.
- Hotopp, K.P. 2002. Land snails and soil calcium in central Appalachian Mountain forest. *Southeastern Naturalist*, 1, 27–44. doi:[10.1656/1528-7092\(2002\)001\[0027:LSA\]2.0.CO;2](https://doi.org/10.1656/1528-7092(2002)001[0027:LSA]2.0.CO;2).
- Hubricht, L. 1964. The land snails of Georgia. *Sterkiana*, 16, 5–10.
- Hubricht, L. 1985. The distributions of the native land mollusks of the eastern United States. *Fieldiana, Zoology*, 24, 1–191. doi:[10.5962/bhl.title.3329](https://doi.org/10.5962/bhl.title.3329).
- Joe, S.M. and C.C. Daehler. 2008. Invasive slugs as under-appreciated obstacles to rare plant restoration: evidence from the Hawaiian Islands. *Biological Invasions*, 10, 245–255. doi:[10.1007/s10530-007-9126-9](https://doi.org/10.1007/s10530-007-9126-9).
- Jones, C.G. and M. Shachak. 1990. Fertilization of the desert soil by rock-eating snails. *Nature*, 346, 839. doi:[10.1038/346839a0](https://doi.org/10.1038/346839a0).
- LaPierre, K.J, W.S. Harpole, and K.N. Suding. 2010. Strong feeding preference of an exotic generalist herbivore for an exotic forb: a case of invasional antagonism. *Biological Invasions*, 12(9), 3025–3031. doi:[10.1007/s10530-010-9693-z](https://doi.org/10.1007/s10530-010-9693-z).
- Lydeard, C., R.H. Cowie, W.F. Ponder, A.E. Bogan, P. Bouchet, S.A. Clark, K.S. Cummings, et al. 2004. The global decline of nonmarine mollusks. *BioScience*, 54(4), 321–330. doi:[10.1641/0006-3568\(2004\)054\[0321:TGDONM\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0321:TGDONM]2.0.CO;2).
- Mason, C.F. 1974. Chapter 17: Mollusca. *In*: C.H. Dickinson, C.H., and G.F.J. Pugh, eds. *Biology of Plant Litter Decomposition*. Academic Press, pp. 564–583.
- Meyer, W.M. III and R.H. Cowie. 2010. Feeding preferences of two predatory snails introduced to Hawaii and their conservation implications. *Malacologia*, 53, 135–144. doi:[10.4002/040.053.0106](https://doi.org/10.4002/040.053.0106).

- Minton, R.L and K.E. Perez. 2010. Analysis of museum records highlights unprotected land snail diversity in Alabama. *American Malacologists Bulletin*, 28, 91–95. doi:[10.4003/006.028.0224](https://doi.org/10.4003/006.028.0224).
- Nichols, B.J. and K.R. Langdon. 2007. The smokies All Taxa Biodiversity Inventory: history and progress. *Southeastern Naturalist*. Special Issue, 1, 27–34. doi:[10.1656/1528-7092\(2007\)6\[27:TSATBI\]2.0.CO;2](https://doi.org/10.1656/1528-7092(2007)6[27:TSATBI]2.0.CO;2).
- Niemiller M.L, D.B. Fenolio, and K.S. Zigler. 2012. The obligate cave fauna of Georgia. *Bulletin of the Georgia Speleological Survey*, 6–12.
- Perez, K.E., N. Defreitas, J. Slapcinsky, R.L. Minton, F.E. Anderson, and T.A. Pierce. 2014. Molecular phylogeny, evolution of shell shape, and DNA barcoding in Polygyridae (Gastropoda: Pulmonata), an endemic North American clade of land snails. *American Malacologists Bulletin*, 32(1), 1–31. doi:[10.4003/006.032.0103](https://doi.org/10.4003/006.032.0103).
- Reeves, W.K., J.B. Jensen, and J.C. Ozier. 2000. New faunal and fungal records from caves in Georgia, USA. *Journal of Cave and Karst Studies*, 62(3), 169–179.
- Schilthuizen, M. and H.A. Rutjes. 2001. Land snail diversity in a square kilometre of tropical rainforest in Sabah, Malaysian Borneo. *Journal of Molluscan Studies*, 67, 417–423. doi:[10.1093/mollus/67.4.417](https://doi.org/10.1093/mollus/67.4.417).
- Slapcinsky, J. 2018. *Vitrea clingmani* Dall in Pilsbry, 1900, a snail endemic to the summits of the Black Mountains and Great Craggy Mountains of North Carolina is now assigned to the genus *Pilsbryna* (Gastropoda: Stylommatophora: Oxychilidae). *The Nautilus*, 132(1), 1–12.
- Smokies species tally. 2016. Discover Life in America. <https://www.dlia.org/smokies-species-tally>. Accessed 17 June 2016.
- Solem, A. 1984. A world model of land snail diversity and abundance. *In*: A. Solem and A.C. van Bruggen, eds. *World-wide Snails: Biogeographical Studies on Non-Marine Mollusca*. Brill Archive, pp. 6–22.
- Turgeon, D.D., J.F. Quinn, A.E. Bogan, E.V. Coan, F.G. Hochberg, W.G. Lyons, P.M. Mikkelsen, et al. 1998. Common and scientific names of aquatic invertebrates from the United States and Canada: mollusks. *American Fisheries Society Special Publication* 26.
- Turke, M., E. Heinze, K. Andreas, S.M. Svendsen, M.M. Gossner, and W.W. Weisser. 2010. Seed consumption and dispersal of ant-dispersed plants by slugs. *Oecologia*, 163, 681–693. doi:[10.1007/s00442-010-1612-6](https://doi.org/10.1007/s00442-010-1612-6).
- US Environmental Protection Agency. 2006. Level III Ecoregions of the Continental United States. National Health and Environmental Effects Research Lab, Corvallis, OR. Map M-1.